THIS PAGE IS INSERTED BY OIPE SCANNING AND IS NOT PART OF THE OFFICIAL RECORD

Best Available Images

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

BLACK BORDERS

TEXT CUT OFF AT TOP, BOTTOM OR SIDES

FADED TEXT

BLURRY OR ILLEGIBLE TEXT

SKEWED/SLANTED IMAGES

COLORED PHOTOS HAVE BEEN RENDERED INTO BLACK AND WHITE

VERY DARK BLACK AND WHITE PHOTOS

UNDECIPHERABLE GRAY SCALE DOCUMENTS

IMAGES ARE THE BEST AVAILABLE COPY. AS RESCANNING WILL NOT CORRECT IMAGES, PLEASE DO NOT REPORT THE IMAGES TO THE PROBLEM IMAGE BOX.

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



| 51) International Patent Classification 5: | | (11) International Publication Number: | WO 94/1434 |
|---|--|--|-------------------------|
| A23L 1/308 | A1 | (43) International Publication Date: | 7 July 1994 (07.07.94 |
| 21) International Application Number: PCI 22) International Filing Date: 24 December 19 | 7/AU93/006 193 (24.12.9 | (AT, BE, CH, DE, DK, ES, FR, | |
| 30) Priority Data: PL 6537 24 December 1992 (24. | 12.92) A | Published With international search report. | |
| 71) Applicant (for all designated States except U MAN FIELDER LIMITED [AU/AU]; Locked Grosvenor Place P.O., Sydney, NSW 2000 (A | l Mail Bag | | |
| 72) Inventors; and 75) Inventors/Applicants (for US only): McNAUG. J. [AU/AU]; 18 Marcella Street, North Eppin. (AU). BROWN, Ian, L. [AU/AU]; 18 Valley. NSW 2227 (AU). GANLY, Robert [AU/AU] Street, Kew, VIC 3101 (AU). | g, NSW 21 Way, Gym | 21 a. | |
| 74) Agent: F.B. RICE & CO.; 28A Montague Str NSW 2041 (AU). | ect, Balma | n, | |
| | | | |
| | | | |
| | | | |
| 54) Title: FOOD COMPOSITIONS INCLUDING R | ESISTANT | STARCH | |
| 54) Title: FOOD COMPOSITIONS INCLUDING R | ESISTANI | STARCH | |
| | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relates than the content. In each case the source of 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relate thanced dietary fibre content. In each case the source f 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relates thanced dietary fibre content. In each case the source f 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relate thanced dietary fibre content. In each case the source f 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relates than the content. In each case the source of 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |
| The invention described in this application relates than the content. In each case the source of 50 % or more, the starch being incorporated into a | es to food o ce of dietar of food comp | ompositions such as breakfast cereals, bread a | naving an amylose conte |

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| AT | Austria | GB | United Kingdom | MR | Mauritania |
|------|--------------------------|-----------|------------------------------|------|--------------------------|
| ΑU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | , NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | rr | Italy | PL. | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgystan | RU | Russian Pederation |
| CF | Central African Republic | KP | Democratic People's Republic | SD | Sudan |
| CG | Congo | | of Korea | SE | Sweden |
| CH | Switzerland | KR | Republic of Korea | SI | Slovenia |
| CI | Côte d'Ivoire | KZ | Kazakhstan | SK | Slovakia |
| CM | Cameroon . | LI | Liechtenstein | SN | Senegal |
| CN | China | ĽK | Sri Lanka | TD | Chad |
| cs | Czechoslovakia | LŪ | Luxembourg | TG | Togo |
| CZ | Czech Republic | LV | Latvia | TJ | Tajikistan |
| DE | Germany | MC | Monaco | TT | Trinidad and Tobago |
| · DK | Denmark | MD | Republic of Moldova | UA | Ukraine |
| ES | Spain | MG | Madagascar | US | United States of America |
| FI | Finland | ML | Mali | UZ | Uzbekistan |
| FR | France | MN | Mongolia | VN | Viet Nam |
| GA | Gabon | | _ | | |

PCT/AU93/00684 WO 94/14342

FOOD COMPOSITIONS INCLUDING RESISTANT STARCH TECHNICAL FIELD

This invention relates to food composition which include resistant starch as a source of dietary fibre and 5 in particular to food compositions which include grain or parts thereof and starch derived from the grain which is high in amylose.

BACKGROUND ART

30

It has been recognised that a balanced diet must 10 include an adequate level of dietary fibre. There are many natural food sources of dietary fibre of which cereals, particularly bran, are recognized as a good source.

However, it would seem that a high proportion of 15 diets of people of the developed countries contain an inadequate level of dietary fibre. An inadequate level of dietary fibre has been linked with a number of diseases which may be broadly classified as metabolic and gastrointestinal. Examples of these diseases are diabetes mellitus, diverticular disease and colonic cancer.

It has also be found that some forms of dietary fibre are useful in lowering blood cholesterol.

One approach to providing adequate levels of dietary fibre has been to produce processed foods which are high 25 in dietary fibre content. Such foods include breakfast cereals, snack bars, bread and the like. Typically bran, outer tissues of the kernel, including pericarp from a variety of cereal sources including wheat, maize, oats and other plant extracts have been used as the source of fibre.

More recently, resistant starches have been recognized as a potential source of dietary fibre. Resistant starches are starches that are highly resistant to hydration and which when ingested pass through the upper regions of the gastrointestinal tract largely 35 unchanged.

- 2 -

To date the only natural sources of resistant starch at a significant level are green bananas and raw potatoes. Such sources do, however, present substantial problems in formulating satisfactory food products, in particular low gelatinization temperatures which are typically 60-80°C.

It is also possible to produce resistant starch by extensive processing involving repeated cooking and cooling of starch pastes.

10 In copending International patent application
No. PCT/AU93/00389 filed 30 July 1993 entitled "High
Amylose Starch and Resistant Starch Fractions" there is
disclosed a hybrid maize seed which is capable of
producing a starch having an amylose content of more than
15 about 80%. That application further discloses a maize
starch derived from such seed having a amylose content of
more than about 80%. This starch was designated as "High
Amylose Starch" in view of the surprising high amylose
content. A number of examples of compositions including
20 high amylose starch were given which included a number of
food applications.

Surprisingly, it has now been found that the aforementioned high amylose starch is high in dietary fibre content and is a resistant starch. Moreover, it has been found that such starches may be advantageously incorporated into food products to achieve enhanced levels of dietary fibre content. Additionally, these starches have relatively high gelatinization termperatures, typically 160-170°C.

30 SUMMARY OF THE INVENTION

Accordingly, the present invention consists in a food composition having an enhanced dietary fibre content, characterised in that the dietary fibre is derived from a starch having an amylose content of at least about 50% or if a rice starch, at least 27% and/or from a grain or parts thereof, the starch content of which has an amylose

PCT/AU93/00684 WO 94/14342

- 3 -

content of at least about 50% or if a rice starch, at least about 27%.

DISCLOSURE OF INVENTION

15

For the purpose of the description that follows, "high amylose" means an amylose content (dsb) of 50% or more, preferably 70% or more, most preferably 80% or more. Particularly preferred amylose contents are 85% or more and 90% or more. For a rice starch, an amylose content of 27% or more is considered to be a "high amylose 10 starch". Note that the method used to determined amylose content is that described in application No. PCT/AU93/00389, the contents of which are incorporated herein by way of reference.

Whilst the resistant nature of the starches of this invention are not fully understood, it is suspected that the molecular arrangement of the glucan polymers, including crystalline structures, and the association of these polymers with other substances, including free fatty acids, may be responsible for imparting the resistant 20 characteristic.

In the accompanying Figure 1, there is shown a graph of total dietary fibre versus amylose content of a number of samples representing three maize varieties-regular maize, waxy maize and high amylose maize. It will be seen 25 from this graph that all the regular maize and waxy maize varieties were substantially nil in dietary fibre content, whereas all of the high amylose samples were found to have a measurable dietary fibre content. Based on the fact that the regular maize samples were found to have an 30 amylose content of about 28% whilst the waxy maize was nil, the finding that at an amylose content of greater than about 50% was associated with the presence of dietary fibre is somewhat surprising. It is also surprising that the dietary fibre content increases to such an extent 35 relative to increasing amylose content.

- 4 -

In Table 1 set out below, it will be seen that when resistant starch is determined by two methods, the content for the high amylose starches, particularly high amylose 80, is substantially greater than regular maize starch.

-5

35

TABLE 1
TOTAL DIETARY FIBRE AND RESISTANT STARCH LEVELS
IN MAIZE STARCH

| | | III IMILUD DIINO. | | |
|----|------------------------------|----------------------------------|--------------------------------|------|
| 10 | | Total Insoluble Dietary Fibre | Resistant Starch | |
| 15 | | (% dsb) | Muir et al McCle (% dsb) (% | - |
| | Waxy Maize Starch 8/91 | ⟨0.2 | 0.7 ± 0.5 | 5.8 |
| 20 | Regular Maize Starch 7/91 | (0.2 | 1.3 ± 0.9 | -0.4 |
| 25 | High Amylose 5 7/91 | 8.0 | 8.2 <u>+</u> 0.4 | 7.5 |
| | High Amylose 8 4/91 | 21.2 | - | 11.9 |
| 30 | High Amylose 8 | 0 31.8 | 20.0 <u>+</u> 1.8 | 18.1 |

Muir et al "Measurement of Resistant Starch factors affecting starch escaping digestion in vitro" Am. J. Clin. Nutr. $\underline{56}$, 123-127 (1992)

McCleary et al "A rapid procedure for total starch measurement in cereal grains and products" Proc. 42nd RACI Cereal Chem. Conf. Christchurch, NZ (1992) Ed. VJ Humphrey-Taylor pp304-312.

- 5 -

The high amylose starch of the invention may be incorporated into a wide range of food products. A non-exhaustive list of such products include: pasta, noodles, instant noodles, breakfast cereals, baked goods such as bread, biscuits, crackers and the like, snack foods, cheese and other dairy based products.

Incorporation of the high amylose starch of the invention into food products will be at a level consistent with the required level of dietary fibre in the finished 10 product. A broadly useful level of incorporation generally lies in the range of from about 5-60%.

It will be appreciated that in some products, high amylose content whole grain or grits derived from the grain may be incorporated alone or in combination with high amylose starch.

Furthermore, the high amylose starch of the invention may be used as a partial replacement for the flour incorporated in products such as bread.

Whilst the dietary fibre and resistant starch results

20 shown respectively in Fig 1 and Table 1 are directed to
maize starch, it should be appreciated that this invention
is not so-limited. For example, high amylose wheat, high
amylose rice, high amylose pea and high amylose barley
starch may also be utilized in the food compositions of

25 this invention.

MODES FOR CARRYING OUT THE INVENTION

In order to better understand the nature of this invention, a number of examples will now be described. BREAKFAST CEREALS

The high amylose starch of the invention can be used to advantage in a variety of ready-to-eat breakfast cereals. These include flaked cereals, extruded flake cereals, extruded gun-puffed cereals, extruded and other shredded cereals, oven-puffed cereals, granola cereals and extruded expanded cereals.

- 6 -

(1) Flaked Cereals

While still in grit form A-958 (Hi-maize - high amylose starch 80%, available from Starch Australasia Limited) can be processed to make a cornflake by using 5 higher than conventional levels of water addition and longer cooking times. The final toasted product (formulation 2571/1) has a light colour and is significantly higher in dietary fibre content as is shown in Table 2.

10 (2) Extruded Flake

An extruded flake product was made by rotary cooking, cold form pelleting, flaking and toasting according to formulation 2562/1. It will be seen from Table 2 that 2562/1 had a significantly higher dietary fibre content as compared with the control.

Formulation: Extruded Flake

| | | <u>2562/1</u> | <u>Control</u> |
|----|----------------------|---------------|----------------|
| | | * | ક |
| | Maize Polenta | 45.0 | 45.0 |
| 20 | Starch A958 | 42.0 | _ |
| | Regular Maize Starch | - | 42.0 |
| | Sugar | 8.7 | 8.7 |
| | Salt | 1.9 | 1.9 |
| | Malt | 2.5 | 2.5 |
| 25 | | | |
| | | 100.0 | 100.0 |

Formulations 2459/2(b) and 2459/2(c) were prepared as 2562/1. However, 2459/2(b) were tempered overnight while 30 for 2459/2(c), the flakes were dried out at $70^{\circ}C$ for 30 minutes prior to flaking.

- 7 -

| Formulation: | Extruded | Flake | (Multigrain) |
|--------------|----------|-------|--------------|
| | | | |

| | | 2459/2(b) | 2459/2(c) |
|----|-------------|-----------|-----------|
| | | 8 | * |
| | Starch A958 | 43.1 | 43.1 |
| 5 | Rice | 11.5 | 11.5 |
| | Oats | 11.5 | 11.5 |
| | Wheat | 20.7 | 20.7 |
| | Sugar | 8.6 | 8.6 |
| | Malt | 2.6 | 2.6 |
| 10 | Salt | 2.0 | 2.0 |
| | | 100.0 | 100.0 |

The textural properties of the resultant flakes were significantly different for each formulation with the 2450/2 forming hollow or pillow-like flakes, whereas 2459/2(c) had a wrinkled, blistered appearance.

The dietary fibre content of the two flake formulations 2459/2(b) and 2459/2(c) are set out in Table 2, from which it will be seen that overnight tempering of the moist pellet containing starch A958 increases the total dietary fibre content of the resultant flake significantly. A further extruded flake product was made according to formulation 2556/1.

Formulation: Extruded Flake

| 25 | | 2556/1 % |
|----|-------------------------|-------------|
| | Wholegrain Calrose Rice | 56.0 |
| | Wheatgerm | 7.0 |
| | Sugar | 6.9 |
| 30 | Salt | 1.5 |
| | Malt | 1.3 |
| | Icing Sugar Mixture | 6.3 |
| | Gluten | 12.6 |
| | Starch A958 | 8.4 |
| 35 | | <u></u> |
| | | 100.0 |

- 8 -

The dietary fibre content for 2446/1 is set out in Table 2.

TABLE 2
Dietary Fibre Content of Breakfast Cereals

| 5 | | | | | |
|----|--------------------|-----------------|----------------|----------------|------------------|
| | <u>Formulation</u> | <u>Moisture</u> | <u>Total</u> | <u>Soluble</u> | <u>Insoluble</u> |
| | | Content | Dietary | Dietary | Dietary |
| | | | <u>Fibre</u> | <u>Fibre</u> | <u>Fibre</u> |
| | | 8 | (% dsb) | (% dsb) | (% dsb) |
| 10 | Cornflake | | | | |
| | (Control) | 4.6 | 4.0 | 1.0 | 2.9 |
| | 2571/1 | 6.2 | 20.7 | 1.2 | 19.5 |
| | 2562/1 | 4.7 | 12.4 | 0.4 | 12.0 |
| 15 | Control | . 6.5 | 3.9 | 0.3 | 3.5 |
| | 2459/2(b) | 4.3 | 17.1 | - | |
| | 2459/2(c) | 5.0 | 15.3 | - | - |
| 20 | 2556/1 | 5.4 | 4.5 | 1.1 | 3.5 |

(3) Wheat Bites

This product is a crunchy "pillow" which can be consumed as a ready-to-eat cereal with milk or eaten piece-wise as a snack.

- 9 -

The formulation of the product is as follows:

| | Ingredient | <pre>% (wet_mix)</pre> |
|----|----------------------|------------------------|
| | Fine wholemeal flour | 36.67 |
| | Castor sugar | 15.35 |
| 5 | Oat flour | 12.80 |
| | Starch A958 | 11.94 |
| | Fine Bran #3 | 8.53 |
| | Vitamin B1 | 0.0051 |
| | Vitamin B2 | 0.0043 |
| 10 | Niacin | 0.0316 |
| • | Iron | 0.0145 |
| | Honey | 1.19 |
| | Water | 13.47 |
| , | | |
| 15 | | 100.00 |

Preparation of the product is as follows:

- (a) premix dry ingredients;
- (b) process cereal mix with honey and water in a twin 20 screw extruder;
 - (c) crimp and cut ropes from extruder to form individual pillows;
 - (d) toast pillows to reduce moisture and develop colour and flavour.

25 The product was found to have excellent crunch after contact with milk for five minutes. It has a distinctive appearance being a pillow with "strand" markings. Total dietary fibre content was found to be 10%, at a moisture content of 2-4%. The product is unusual in that it has a light airy texture with a relatively high level of total dietary fibre.

The cereal products described above were characterized by several physical properties which were quite unexpected. For example, some of the flakes were blistered whilst others were bubbled and appeared as if

- 10 -

"popped". These characteristics are highly desirable and would not have been considered achievable using formulations of this type.

In addition, these cereal products extrude well,

5 display moderate to high expansion, yielding a final
product with a medium to coarse cell structure. The
products are unusual in that they have a light airy
texture whilst containing a relatively high level of total
dietary fibre.

The breakfast cereal products have also been found to have an excellent "bowl life" in that the flakes retained their crunch after contact with milk for five minutes.

Furthermore, during processing and prior to toasting, it was observed that the flakes had a tendency to stick less to each other as compared with the control.

BREAD

A number of bread samples were prepared using a high amylose starch and for comparison a waxy maize starch "MAZACA" trade mark of Starch Australasia Limited. The 20 starches were incorporated into the bread at a 5% and a 10% level of flour replacement.

In Table 2, set out below, are the results obtained for each of the bread samples. These results clearly show that at both the 5% and 10% levels of high amylose starch, the dietary fibre content was substantially more than both the control product and the mazaca based product. Furthermore, it should be noted that the actual level of dietary fibre found was significantly more than would be expected. This suggests that a synergistic effect occurred during baking.

Indeed, it would have been expected that the dietary fibre content would decrease as resistant starch is destroyed during such heating processes.

- 11 -

| | • | TABLE 3 | | |
|----|-----------------------|-----------|---------|-------------|
| | Analysis | Wheat | High | Mazaca |
| | | Starch | Amylose | |
| | | Control | Starch | |
| 5 | | | | |
| | | | | |
| | Moisture % | - | 13.0 | 12.6 |
| | Total dietary fibre | - | 27 | 0 |
| | (TDF) % (DB) | | | |
| 10 | (Resistant starch) | | | |
| | Test baking | | | |
| | Rate of flour | 0 5 10 | 5 10 | 5 10 |
| 15 | replacement (%) | | : | |
| | Notes observation (%) | | 58 58 | 58 58 |
| | Water absorption (%) | | | |
| | Total loaf score (%) | 76 80 76 | 75 73 | 79 71 |
| 20 | Fibre in bread | | | |
| | TDF % (DB) | 4.2 4.0 - | 6.9 8.7 | 4.0 |
| | (approx. expected | | | |
| 25 | *TDF %) | · | 5.2 6.4 | 4.0 |

[&]quot;expected" bread TDF is based on control loaf fibre plus raw starch fibre.

WO 94/14342

- 12 -

A further series of test bakings were done according to the formulations set out below:

| | <u>Ingredients</u> | Batch Nos. | | | | | |
|----|--------------------|------------|-----|-----|-----|-----|-----|
| | | _1_ | _2_ | _3_ | _4_ | _5_ | _6_ |
| 5 | Flour (g) | 1000 | 950 | 900 | 850 | 800 | 750 |
| | Starch A.958 (g) | 0 | 50 | 100 | 150 | 200 | 250 |
| | Gluten (g) | 0 | 10 | 20 | 30 | 40 | 50 |
| | Salt (g) | 18 | 18 | 18 | 18 | 18 | 18 |
| | Improver (g) | 15 | 15 | 15 | 15 | 15 | 15 |
| 10 | Fat (g) | 20 | 20 | 20 | 20 | 20 | 20 |
| | Yeast (g) | 12 | 12 | 12 | 12 | 12 | 12 |
| | Water (mL) | 600 | 610 | 630 | 650 | 670 | 700 |

Each batch was processed in a conventional manner. The total dietary fibre for each batch was found to be:

| | Batch No. | Total Dietary Fibre |
|----|-----------|---------------------|
| | • | (% dsb) |
| | 1 | 5.4 |
| | 2 | 7.1 |
| 20 | 3 | 9.1 |
| | 4 | 10.8 |
| | 5 | 12.8 |
| | 6 | 14.5 |

In addition, it was noted that inclusion of the high 25 amylose starch did not have an adverse effect on crumb colour, whilst increasing levels of high amylose starch led to increasing dough water absorption levels.

Generally it was concluded that the high amylose starch was an excellent source of dietary fibre for inclusion in bread since any detrimental effect on load volume and crumb firmness could be readily overcome by appropriate formulation changes known to those skilled in the art.

It has also been found that gluten free bread which is high in dietary fibre may be produced using the high

- 13 -

amylose starch of the invention. This is important as conventional gluten free bread, which is consumed by sufferers of coeliac disease, is generally low in dietary fibre.

For guidance, up to about 15% may be incorporated into a gluten free bread to yield a dietary fibre content of about 10%.

As well as being able to provide enhanced dietary fibre content in bread, the high amylose starch acts to reduce staling on storage of the bread, by it is thought, reducing the level of amylopectin and helping to maintain the moisture content of the bread.

NOODLES

Typically, up to about 24% of high amylose starch may 15 be used as a flour replacement in spaghetti. This yields a dietary fibre content of up to about 8.7%.

In use, it has been found that spaghetti made from high amylose starch has less of the "speckiness" which is usually associated with the addition of more conventional fibre additivies such as bran to achieve higher dietary fibre content. Furthermore, during cooking, the spaghetti is less sticky and remains all dente for longer.

OTHER FOOD PRODUCTS

It has been found that, in addition to providing a dietary fibre benefit in instant noodles, these products have a greater crispiness as compared with conventional formulated products.

In the formulation of biscuits and crackers, it is possible to overcome the use of a laminator by using the 30 high amylose starch of the invention. This is due to the expansion, popping effect which serves to impart a lighter texture to such products.

INDUSTRIAL APPLICABILITY

From the foregoing description it will be evident that the present invention provides a means for enhancing

- 14 -

the dietary fibre content of a variety of food compositions. Not only does this have obvious nutritional benefits, but it allows for the production of food compositions having physical characteristics not previously known with conventional formulations.

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

5

- 15 -

CLAIMS

1. A food composition having an enhanced dietary fibre content, characterised in that the dietary fibre is derived from a starch having an amylose content of at


- 5 least about 50% or if a rice starch, at least about 27%, and/or from a grain or parts thereof, the starch content of which has an amylose content of at least about 50% or if a rice starch, at least about 27%.
- 2. A food composition as in claim 1 wherein the amylose 10 content of the starch is 55% or more.
 - 3. A food composition as in claim 2 wherein the amylose content of the starch is 70% or more.
 - 4. A food composition as in claim 3 wherein the amylose content of the starch is 80% or more.
- 15 5. A food composition as in claim 4 wherein the amylose content of the starch is 85% or more.
 - 6. A food composition as in claim 5 wherein the amylose content of the starch is 90% or more.
- A food composition as in any one of claims 1 to 6
 wherein the starch is a wheat, maize, barley, pea or rice starch and/or the grain or parts thereof is wheat, maize, barley, pea or rice.
 - 8. A food composition as in claim 7 wherein the starch is maize starch and/or the grain or parts thereof is maize.
- 25 9. A food composition as in any one of claims 1 to 8 wherein the starch and/or the grain or parts thereof are included in an amount of from about 5-60% to give a dietary fibre content of about 1.5-20%.
- 10. A food composition as in claim 9 comprising noodles 30 wherein the starch is in an amount of up to about 24%.
 - 11. A food composition as in claim 9 comprising bread wherein the starch is in an amount of from about 5-25%.
- 12. A food composition as in claim 11 wherein the bread is gluten free and includes the starch in an amount up to 35 about 15%.

- 16 -

13. A food composition as in claim 9 comprising a breakfast cereal.

- 14. A food composition as in claim 13 wherein the breakfast cereal is selected from the group consisting of
- 5 flaked cereals and extruded flakes.
 15. A food composition as in claim 14 wherein the
 - breakfast cereal is bubbled, popped or blistered in appearance.
- 16. A food composition as in claim 15 wherein the dietary 10 fibre content is about 4.5% or more.
 - 17. A food composition as in claim 16 wherein the dietary fibre content is about 12.4% or more.
 - 18. A food composition as in claim 17 wherein the dietary fibre content is about 15.3% or more.
- 15 19. A food composition as in claim 18 wherein moist pellets produced during formation of the flakes are tempered overnight, such that the dietary fibre content thereof is about 17.5% or more.
- 20. A food composition as in claim 17 wherein the dietary 20 fibre content is about 20.7% or more.
 - 21. A food composition as in claim 15 wherein the breakfast cereal is in the form of "pillows" having a light, airy texture.
- 22. A food composition as in claim 21 wherein the
- 25 breakfast cereal has a dietary fibre content of about 10%.

TOTAL DIETARY FIBRE (% d.s.b.)



■ Regular Maize

Waxy Maize

High Amylose Maize

FIG 1

SUBSTITUTE SHEET (Rule 26)

| | CLASSIFICATION OF SUBJECT MATTER CL. 5 A23L 1/308 | | | | | | | |
|--|--|--|-----------------------|--|--|--|--|--|
| According to International Patent Classification (IPC) or to both national classification and IPC | | | | | | | | |
| В. | B. FIELDS SEARCHED | | | | | | | |
| Minimum documentation searched (classification system followed by classification symbols) IPC A23L 1/308; A23L 1/30 (up to and including 1983 | | | | | | | | |
| Documentation AU:IPC as a | on searched other than minimum documentation to above | the extent that such documents are included in | n the fields searched | | | | | |
| Electronic data base consulted during the international search (name of data base, and where practicable, search terms used) DERWENT: DIET: () FIBRE, AMYLOSE OR STARCH CHEMICAL ABSTRACTS: DIET: () FIBRE, AMYLOSE OR STARCH | | | | | | | | |
| C. | DOCUMENTS CONSIDERED TO BE RELEVA | ANT | | | | | | |
| Category* | Citation of document, with indication, where a | appropriate, of the relevant passages | Relevant to Claim No. | | | | | |
| x | AU-A-25124/92 (A.E STALEY MANUFA 4 March 1993 (04.03.93) Claim 16 | 1-22 | | | | | | |
| x | AU-B-61403/90 (628218) (AMERICAN M. 22 February 1991 (22.02.91) Whole document | | 1-22 | | | | | |
| x | AU-B-60630/90 (628206) (AMERICAN M. 22 February 1991 (22.02.91) Claim 2 | AIZE-PRODUCTS COMPANY) | 1-22 | | | | | |
| X Further in the | Further documents are listed in the continuation of Box C. | | | | | | | |
| * Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance earlier document but published on or after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention document of particular relevance; the claimed invention cannot be considered novel or cannot be or which is cited to establish the publication date of another citation or other special reason (as specified) document referring to an oral disclosure, use, exhibition or other means document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention cannot be considered novel or cannot be considered to involve an invention cannot be considered to invention cann | | | | | | | | |
| | tual completion of the international search | Date of mailing of the international search | · \ | | | | | |
| | 094 (25.03.94) | | 03,94) | | | | | |
| AUSTRALIA PO BOX 200 WODEN AC AUSTRALIA | | J. BODEGRAVEN | ŕ | | | | | |
| Facsimile No. | Facsimile No. 06 2853929 Telephone No. (06) 2832281 | | | | | | | |

| egory* | Citation of document, with indication, where appropriate of the relevant passages | Relevant to Claim No. |
|--------|---|-----------------------|
| | FR,A,2518372 (SOCIETE D'HYGIENE NUTRITIONELLE (SHN)) 24 June 1983 (24.06.83) | |
| X | Claim 3, examples | 1-22 |
| | Cereal Chemistry 68(6) 1991 pages 589-596 Szczodrak J. and Pomeranz Y. "Starch and Enzyme-Resistant Starch from High-Amylose Barley" | 1-22 |
| X | Whole document | 1-22 |
| | AU-B45616/89 (625542) (IOWA STATE UNIVERSITY RESEARCH FOUNDATION, INC.) 31 May 1990 (31.05.90) | |
| x | Whole document | 1-22 |
| | | |
| | | |
| | • | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | · | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |

END OF ANNEX

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

| | Patent Document Cited in Search Report | i, | | | Patent Family | Member | | |
|----|--|----------|-------------------|----|---------------|--------|---------|---|
| AU | 25124/92 | EP | 529891 | IL | 102829 | wo | 9303628 | |
| ΑU | 61403/90 | EP | 485436 | US | 4981709 | wo | 9101092 | |
| AU | 60630/90 | EP | 482094 | US | 5094872 | | | |
| AU | 45616/89 | EP US | 372358 5004864 | HU | 896217 | HU | 63298 | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | : | | | | | | |
| | | ٠ | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | • | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | • |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |
| | | | | | | | | |